

NEW SPECIES

'*Phoce* *massiliensis*' a new bacterial species isolated from the human gut

S. Ndongo¹, J.-C. Lagier¹, P.-E. Fournier¹, D. Raoult^{1,2} and S. Khelaifia¹

1) Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, Aix-Marseille Université, Marseille, France and 2) Special Infectious Agents Unit, King Fahd Medical Research Centre, King Abdulaziz University, Jeddah, Saudi Arabia

Abstract

We present here the main characteristics of '*Phoce* *massiliensis*' strain Marseille-P2769^T (CSUR P2769), which is a new genus isolated from the stool of a 45-year-old patient.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: Culturomics, Genomics, Human-gut, '*Phoce* *massiliensis*', taxonomy

Original Submission: 1 June 2016; **Revised Submission:** 9 June 2016; **Accepted:** 9 June 2016

Article published online: 15 June 2016

Corresponding author: S. Khelaifia, Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, Aix-Marseille Université, Marseille, France
E-mail: khelaifia_saber@yahoo.fr

In March 2016, as a part of our culturomics studies [1], we isolated from the stool sample of a 45-year-old patient hospitalized for the treatment of a melanoma, a bacterial strain that could not be identified by our systematic matrix-assisted laser desorption–ionization time-of-flight mass spectrometry (MALDI-TOF-MS) screening on a MicroFlex spectrometer (Bruker Daltonics, Bremen, Germany) [2]. This study was approved by the local committee of the IFR48 (Marseille, France) under agreement number 09-022 and the consent of the patient was obtained. The initial growth was observed after the stool sample had been pre-incubated in a blood bottle culture (Becton-Dickinson, Pont de Claix, France), with the addition of 5% sheep rumen for 5 days. Several colonies appeared after 3 days of cultivation on 5% sheep blood agar (bioMérieux, Marcy l'Etoile, France) in anaerobic conditions generated by AnaeroGenTM (bioMérieux). Agar-grown colonies ranged from 1 to 2 mm in diameter and were beige and circular. Bacterial cells were Gram-negative bacilli, strictly anaerobic, motile and non-spore-forming (0.4–0.5 × 2.5–3 µm). Strain Marseille-P2769 was catalase and oxidase negative. As previously described, we sequenced the

complete 16S rRNA gene using a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France) with the universal primers FDI and RP2 (Eurogentec, Angers, France) [3]. Strain Marseille-P2769 exhibited a 92% sequence identity with *Anaerotruncus colihominis* strain JM4-15^T (GenBank accession number KR364734), the phylogenetically closest species with standing in nomenclature. Indeed, with a similarity <95%, strain Marseille-P2769 was putatively classified as a member of a new genus within the family Ruminococcaceae in the Firmicutes phylum [4].

Anaerotruncus colihominis was isolated from human faeces and was described in 2004 [5]. It was obligatory anaerobic, Gram-positive, non-spore-forming and cells were rods (0.5–2.5 µm). Catalase was not produced.

Strain Marseille-P2769 exhibited a 16S rRNA sequence divergence >5% with its phylogenetically closest species with standing in nomenclature [6], so we propose the creation of the new genus '*Phoce*' gen. nov., (pho.ce.en'sis. L. fem. adj. *phoceensis* referring to Phoce, the Greek name of the city which founded Marseille, where the strain was isolated). Strain Marseille-P2769^T is the type strain of the new species '*Phoce* *massiliensis*' gen. nov., sp. nov (see Fig. 1).

MALDI-TOF-MS spectrum accession number

The MALDI-TOF-MS spectrum of '*Phoce* *massiliensis*' is available at <http://www.mediterranee-infection.com/article.php?leref=256&titre=urms-database>.




FIG. 1. Phylogenetic tree showing the position of 'Phoceia massiliensis' strain Marseille-P2769^T relative to other phylogenetically close members of the family Ruminococcaceae. GenBank accession numbers are indicated in parentheses. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using the maximum-likelihood method within the MEGA software. Numbers at the nodes are percentages of bootstrap values obtained by repeating the analysis 500 times to generate a majority consensus tree. Only values >70% were displayed. The scale bar indicates a 2% nucleotide sequence divergence.

Nucleotide sequence accession number

The 16S rRNA gene sequence of strain Marseille-P2769 was deposited in GenBank under accession number LT558849.

Deposit in a culture collection

Strain Marseille-P2769 was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR, WDCM 875) under number P2769.

Conflict of Interest

The authors have no conflicts of interest to declare.

Acknowledgements

This research was funded by Méditerranée-Infection Foundation.

References

- [1] Lagier J-C, Hugon P, Khelaifa S, Fournier P-E, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev* 2015;28: 237–64.
- [2] Seng P, Abat C, Rolain JM, Colson P, Lagier J-C, Gouriet F, et al. Identification of rare pathogenic bacteria in a clinical microbiology laboratory: impact of matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J Clin Microbiol* 2013;51:2182–94.
- [3] Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol* 2000;38: 3623–30.
- [4] Euzéby J. Validation List No. 132. *Int J Syst Evol Microbiol* 2010;60: 469–72.
- [5] Lawson PA, Song Y, Liu C, Molitoris DR, Vaisanen M-L, Collins MD, et al. *Anaerotruncus colihominis* gen. nov., sp. nov., from human faeces. *Int J Syst Evol Microbiol* 2004;54:413–7.
- [6] Kim M, Oh HS, Park SC, Chun J. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* 2014;64:346–51.